

Supplementary Materials for **Specific and Nonhepatotoxic Degradation of Nuclear Hepatitis B Virus cccDNA**

Julie Lucifora, Yuchen Xia, Florian Reisinger, Ke Zhang, Daniela Stadler, Xiaoming Cheng,
Martin F. Sprinzl, Herwig Koppensteiner, Zuzanna Makowska, Tassilo Volz,
Caroline Remouchamps, Wen-Min Chou, Wolfgang E. Thasler, Norbert Hüser, David Durantel,
T. Jake Liang, Carsten Münk, Markus H. Heim, Jeffrey L. Browning, Emmanuel Dejardin,
Maura Dandri, Michael Schindler, Mathias Heikenwalder,* Ulrike Protzer*

*Corresponding author. E-mail: protzer@tum.de (U.P.); heikenwaelder@helmholtz-muenchen.de (M.H.)

Published 20 February 2014 on *Science Express*
DOI: 10.1126/science.1243462

This PDF file includes:

Materials and Methods

Figs. S1 to S21

Table S1

References

Supplementary Materials

Materials and Methods

HBV inocula, cell cultures, HBV infection and treatments

HBV inocula were prepared as described (42) Shortly, HBV was concentrated from the supernatant of HepG2.2.15 cells using centrifugal filter devices (Centricon Plus-70, Biomax 100.000, Millipore Corp., Bedford, MA) and titered by HBV-DNA dot blot analysis after sedimentation into a CsCl density-gradient to determine enveloped DNA-containing viral particles (vp). Only inocula reaching a titer between 3×10^9 and 3×10^{10} vp/mL were used. Immediately after collection, the virus stock was aliquoted and stored at -80°C until used for infection. Primary Human Hepatocytes (PHH) were isolated from surgical liver resections, cultured and infected with HBV as described (43, 44). Tissue samples and annotated data were obtained and experimental procedures were performed within the framework of the non profit foundation Human Tissue & Cell Research (HTCR), including the informed patient's consent (45). HepG2-H1.3, HepaRG and HepaRG-tA cell culture, differentiation and HBV infection (at a multiplicity of infection of 600 vp/cells) were also performed as described (13,46). HepaRG-tA-VifAU1 cell line was obtained after transduction of HepaRG-tA cells with a lentivirus expressing the HIV-Vif gene under tetracyclin regulated promoter (Plvx-Tight-Puro lentiviral vector, Clontech, Saint-Germain-en-Laye, France) and selection using 0.25 µg/ml puromycin. A3A or A3B overexpression in HepG2H1.3 cells was obtained by transfection of pLenti6.3, pLenti6.3-A3A, pTR600 or pTR600-A3B, respectively, using “Lipofectamine 2000 Reagent” (Invitrogen). No selection took place before experiment start. HepG2-H1.3-A3A cell line was obtained after lentiviral transduction of A3A into HepG2-H1.3 cells (47) and selection using 5µg/ml blasticidin. HepaRG cells were transfected with episomal replicon construct pEpi-H1.3 using “Lipofectamine® 2000 Reagent” (Invitrogen). pEpi-H1.3 was derived by cloning a 1.3-fold HBV genome into the extrachromosomal mammalian replicon plasmid pEpi-eGFP (kindly provided by H.J. Lipps, Witten-Herdecke, Germany). Two days after transfection, selection with 600 µg/ml geneticin was started and kept until cells expressed GFP and reached confluency and expressed GFP. HepaRG+pEpi-H1.3 cells were differentiated for 6 weeks to allow replication of HBV and establishment of HBV cccDNA by nuclear reimport of HBV rcDNA containing

capsids. Lamivudine and entecavir were used at 0.5 μ M (5-fold and 1000-fold IC 50, respectively), IFN- α (Roferon-A, Roche) at 1000 U/mL and BS1, CBE11 and hu-IgG at 0.5 μ g/mL unless otherwise indicated.

Analysis of HBV replication intermediates

HBeAg was determined using commercial immunoassays (Siemens Molecular Diagnostics, Marburg, for cell culture supernatants; Abbott Laboratories, Wiesbaden, Germany for mouse sera). Total DNA was purified from infected cells using a “Tissue kit” (Macherey Nagel, Düren, Germany). Total RNA was extracted from infected cells (NucleoSpin® RNA II kit, Macherey Nagel) and transcribed into cDNA using SuperScript III reverse transcriptase (Invitrogen). HBV-DNA, cccDNA and pgRNA were detected using specific PCR primers as described (48,17). pEpi-H1.3 was quantified using Kan-L/-R primers (see primer list). Real-time PCRs (qPCR) were performed using the LightCycler™ system (Roche Diagnostics) and analyzed using the second derivative maximum method that includes both normalization to the reference gene (*PrnP* for cccDNA and pEpi-H1.3 and GAPDH for pgRNA) and to primer efficiency (47,49). cccDNA specific primers span the gap and the nick in the rc form of the HBV genome and were designed not to detect rcDNA or a linear 1.1 or 1.3-fold HBV genome. Using optimized PCR conditions on the Light Cycler instrument (95°C for 10'', 45 cycles at 95°C for 15'', 60°C for 5'', 72°C for 45'', and detection at 88°C for 2'' after each cycle), we determined the specificity to amplify cccDNA over rcDNA to be 10³ to 1. For Southern Blot analyses, cccDNA was extracted from HBV-infected cells using the KCl protein precipitation method, separated through 0.8 % agarose gel, blotted onto nylon membrane and hybridized with ³²P HBV-DNA probe (50, 51). Quantification of cccDNA by qPCR was confirmed by quantification from Southern blotting as exemplarily shown in Figure S3B.

3D-PCR

PCR products from cccDNA-specific qPCR were diluted 1/20 to 1/50 and used as templates for nested PCR using HBxin fw and HBxin rev primers (see primer list) and decreasing denaturation temperatures as indicated. PCR products were purified by gel extraction (GeneJET Gel Extraction Kit, Fermentas, Fisher Scientific - Germany, Schwerte, Germany) and cloned by TA cloning (TA Cloning® Kit, Invitrogen). DNA from individual clones was purified (GeneJET

Plasmid Miniprep Kit, Fermentas, Fisher Scientific - Germany GmbH) and sequenced using primers hybridizing to the 5' and 3' plasmid backbone.

AP site quantification and repair

AP site quantification was performed with OxiSelect™ Oxidative DNA Damage Quantitation Kit (AP sites) (STA-324, Cell Biolabs, San Diego, CA, USA) according to the manufacturer's instruction using 0.5 µg of purified DNA (100 µg/mL). 1µg of total DNA were digested with 10 units of apurinic apyrimidinic endonuclease redox effector factor-1 (APE1; M0282L, New England Biolabs, Inc., Ipswich, MA, USA) or 1 µl PreCR Repair Mix (M0309L, New England Biolabs, Inc.).

qRT-PCR

For qRT-PCR RNA was extracted using commercial kits (RNeasy-kit from Qiagen, Hilden, Germany or NucleoSpin® RNA II kit, Macherey Nagel) and transcribed into cDNA with the Quantitect Reverse Transcription kit (Qiagen) or with SuperScript III reverse transcriptase (Invitrogen) according to the manufacturer's instructions. Expression levels were quantified by qRT-PCR performed on a 7900 HT qRT-PCR system (Applied Biosystems, Life Technologies Darmstadt, Germany) using the $\Delta\Delta CT$ method or on a LightCycler™ system (Roche Diagnostics) and analyzed using the second derivative maximum method that includes both normalization to the reference genes and primer efficiency. Relative mRNA levels of all target genes were normalized to at least two house-keeping genes (HPRT; RHOT2; TBP; GAPDH) levels.

siRNA and antibodies

siRNAs were obtained from Thermo Fisher Scientific Dharmacon® ([Lafayette, CO, USA](#)) : control siRNA (D-001910-10-05 Acell non-targeting pool 5nmol), siRNA against A3B (E-017322-00-0005, Accell SMARTpool, Human A3B (9582), 5 nmol) or siRNA against A3A (#E-017432-00, Accell SMARTpool). For Western blot analyses, immunostaining and immunoprecipitation, the following primary antibodies were used: polyclonal anti-NFkB2 p100/p52 (#4882, Cell Signaling Technology, Inc, Danvers, MA, USA), anti-phospho-RelA (#RB-1638-P0, Cell Signaling Technology), monoclonal anti-GAPDH (clone 14C10) (Cell

Signaling Technology), anti- β actin (clone 13E5) (Cell Signaling Technology), polyclonal anti-LT β R (#sc-8375, Santa Cruz Biotechnology, Inc, Heidelberg, Germany), anti-phospho-NIK (#4994, Santa Cruz Biotechnology), anti-TNFRI (#ab19139, Abcam, Cambridge, MA, USA), rabbit anti-HBVcore antiserum H800 (kindly provided by Heinz Schaller, Heidelberg, Germany), rabbit polyclonal anti-HBVcore (B0586, Dako, Hamburg, Germany), mouse monoclonal anti-HBVcore (F8.27.2, kindly provided by Dr. Marie-Anne Petit), anti-RelB (#sc-226, Santa Cruz Biotechnology), polyclonal anti-A3A (#AP31973PU-N, Acris Antibodies GmbH, Herford, Germany), polyclonal anti-A3F (#H00200316-A01; Abnova, Aachen, Germany), polyclonal anti-A3G (#AP23049PU-N, Acris Antibodies GmbH), polyclonal anti-AU1 (#A190-125A-2, Bethyl Laboratories, Inc., Montgomery, TX, USA), polyclonal anti-A3B (#sc-130955, Santa Cruz Biotechnology Inc), anti-Flag (F7425, Sigma), anti-TBP polyclonal antibody (H00006908-A01, Bethyl, USA), polyclonal anti-histone H3 (#ab1791, Abcam).

Deep sequencing

Target genes were amplified from total DNA by PCR using primers (see table below) designed to span around 2000bp of genomic sequence containing at least one exon and one intron. PCR reaction was performed with the Advantage®-HF2 PCR-kit (Clontech). Subsequently, amplicons were separated by a 0.8% agarose gel and purified using the S.N.A.P.™ UV-free gel purification kit (Life Technologies). Purified amplicons were submitted for deep sequencing to a commercial provider (GATC Biotech, Konstanz, Germany) and analyzed using Hiseq2000 sequencing system (Illumina, San Diego, CA, USA). Sequencing data were analyzed with CLC Genomics Workbench 6.01 (CLC bio, Aarhus, Denmark).

Nr.	Oligoname	Sequence (5'->3')
1	B2M fwd	CTGCCGTGTGAACCATGTGA
2	B2M rev	GCTACCTGTGGAGCAACCTG
3	CCL17 fwd	ACACAGAGACTCCCTCCTGG
4	CCL17 rev	GCTCCAGTTTCAGACAAGGGG
5	CXCL10 fwd	AGCAGAGGAACCTCCAGTCT
6	CXCL10 rev	TGTGGTCCATCCTTGGAAAGC
7	CXCL13 fwd	CGACATCTCTGCTTCTCATGC
8	CXCL13 rev	AGACTGAGCTCTTGACAC
9	HMBS fwd	GGGCTCAGTGTCTGGTTAC
10	HMBS rev	CCCAGAGCCCTCTAGACCTT

11	IL-6 fwd	GAGGAGACTTGCCTGGTGAAA
12	IL-6 rev	GCCCATGCTACATTGCCG
13	MxA fwd	TCCATGACTCGCAGAGAGGA
14	MxA rev	TCCACTGTAGAACACGCCAC
15	MYC fwd	CCACCTCCAGCTTGTACCTG
16	MYC rev	GCTGCGTAGTTGTGCTGATG
17	PKR fwd	CTTGCCGTCACGGGGATAAT
18	PKR rev	CTTGGCATTAAAGTCGCTGCC
19	PML fwd	GAAAGCCCCAAGCCAACAGG
20	PML rev	ATTGGCAGGATGGTTGAGGG
21	SRC fwd	TGGAACTTGCCAGGGTTGTT
22	SRC rev	ACCAGGGCAGGAAATGTAGC
23	TBP fwd	TTGGTAGTTGAATCCCGCCC
24	TBP rev	TCACTGTGAAGAGAGCGCAG

Immunofluorescence microscopy

Cells grown on 4-well-glass slide (Lab-Tek II, Fisher Scientific - Germany, Schwerte, Germany) were fixed with 4% paraformaldehyde (PFA) in PBS pH 7.4 for 10 min at room temperature and permeabilized with 0.5% saponin. Slides were blocked in PBS containing 0.5% saponin and 10% serum from the species that the secondary antibody was raised in at room temperature for two hours. After blocking, primary antibodies diluted in PBS with 0.1% saponin and 10% blocking serum were added overnight at 4°C. After extensive washing, slides were incubated with the secondary antibody in PBS with 0.1% saponin and 2% blocking serum for 2 hours at room temperature in the dark. Slides were mounted with Dapi Fluoromount-G (SouthernBiotech, Birmingham, Alabama, USA) and images were captured with a Olympus FV10i confocal Microscope (Olympus, Germany).

Expression profile analyses

Total RNA was amplified and labeled using WT Expression Kit (Ambion, Life Technologies) and Terminal labeling Kit (Affymetrix, Santa Clara, CA, USA) according to the manufacturer's recommendations. The amplified and fragmented, biotinylated complementary RNA was hybridized to Affymetrix Human Gene 1.0 ST Arrays (33297 probe sets) using standard procedures. The experimental setup contained a total of 8 arrays, made up of 2 groups, with each group consisting of 4 biological replicates. Arrays were assessed for quality and robust multi-array average (RMA)-normalized. Quality assessment consisted of RNA degradation plots,

Affymetrix quality control metrics, sample cross-correlation, and probe-level visualizations. Normalization included (separately for each RNA-type data set) background correction, quantile normalization, and probe-level summation by RMA. Data were analyzed for differential gene expression using an empirical Bayes moderated t-test (52), implemented in the Bioconductor package Linear Models for Microarray Data LIMMA. Results were sorted by the adjusted p-value and exported in tab-delimited format. Samples with an up- or downregulation ≥ 2 were re-analysed by qRT-PCR. Microarray data have been submitted to the GEO database (<http://www.ncbi.nlm.nih.gov/geo/>) and have the accession number GSE46667.

Immunoprecipitation assays

Co-Immunoprecipitations were performed using Pierce Co-Immunoprecipitation Kit (Fisher Scientific - Germany, Schwerte, Germany) according to the manufacturer's instructions. Chromatin immunoprecipitation was performed using Pierce Agarose ChIP Kit (#26156, Fisher Scientific) or ChIP kit #ab500 (Abcam, Cambridge, UK). Non-sheared DNA for cccDNA or MNase digested DNA for genomic DNA was analyzed by specific qPCR (GAPDH, P53, SRC, cMyc promoter regions).

FACS-based FRET

For FACS-FRET analysis we used a FACS CantoII Cytometer (BD Bioscience) with 405 nm, 488 nm and 633 nm lasers. Gating strategy and experimental procedures were done as described previously (14). eYFP was excited with 488 nm and detected with a 529/24 filter (Semrock, New York, USA) and eCFP was excited with 405 nm and detected via the 450/40 filter (Semrock). Huh7.5 hepatoma cells transfected with 1.6 μ g DNA per 250.000 cells using Lipofectamine (Life technology) were used for FRET analysis. After 24 h, cells were washed, trypsinized and analyzed by flow cytometry using excitation at 405 nm followed by signal detection with the 529/24 filter. Mock transfected cells, eCFP and eYFP only as well as eCFP/eYFP co-transfected cells served as controls to exclude false positive FRET and subtract background signals. An eCFP-eYFP fusion construct was used as positive control. FRET constructs used within this study were all fused to eCFP/eYFP via their C-terminus using the *NheI-AgeI* restriction sites as previously described (53). PCR derived inserts were sequenced to confirm their identity.

Proximity ligation assay (PLA)

For PLA, cells were fixed with 1 % PFA, washed and permeabilized for 20 min with 1 % Saponin in PBS and blocked for 45 min with PLA blocking solution. Polyclonal rabbit anti-HBVcore and goat anti-A3A antibodies were incubated at a concentration of 1:100 in 1 % BSA for 2 h at RT. Secondary antibody probe incubation, ligation and amplification reactions were done according to the manufacturer's instructions (Duolink, Sigma Aldrich). Spinning disc microscopy with a Nikon Ti Eclipse microscope equipped with a Perkin Elmer UltraViewVox System (Yokogawa CSU-X1) was done for sample imaging. Images were analyzed with the Velocity 6.2 software package (Perkin Elmer) and the implemented automated spot counting was used to assess the number of spots per cropped cell. Statistical analysis was done with GraphPad Prism 5.0 and a one-way analysis of variances (ANOVA).

Modeling of HBc/A3A interaction

A three-dimensional structure of HBV core protein was generated by automatic modeling mode of SWISS-MODEL server (swissmodel.expasy.org) (54). The crystal structure of the human hepatitis B virus capsid (PDB code 1QGT, chains D) (55) was adapted as a template to perform the homology-modeling. In silico docking of A3A (PDB code 2M65) (56) and modeled HBVcore was carried out by using Hex protein docking server (<http://hexserver.loria.fr/>) (57). The result was displayed and analyzed by Rasmol 2.7.5 (58).

HBV-transgenic mice and tissue analysis

6 months old, male hepatitis B virus transgenic mice (C57BL/6, strain HBV1.3.32) (59) received murine lymphotoxin-receptor agonizing (ACH6) or isotype control antibodies intra-peritoneally (2 mg/kg, body weight) three times a week. Animals were maintained under specified pathogen-free conditions. Experiments were performed in accordance to the German legislation governing animal studies and the Principles of Laboratory Animal Care guidelines, NIH (55.1-1-54-2531.3-27-08). Mice were sacrificed at indicated time points and serum as well as liver tissues were obtained. Murine livers were fixed in phosphate buffered saline containing 4% PFA for 24 hours. Fixed liver specimens were embedded in paraffin and 2 µm sections were mounted on glass slides. Immuno-histochemical stainings were performed with cleaved caspase 3 (Cell Signaling), HBVcore (Diagnostic Biosystems, Pleasanton, CA, USA), RelA (NeoMarkers / Lab vision

corporation, Fremont, CA, USA) and RelB (Santa Cruz Biotechnology Inc.) specific antibodies employing an automated BOND-MAX™ platform (Leica Biosystems, Wetzlar, Germany). Positively stained cells were counted in five independent high-power fields and extrapolated to numbers per square millimeter tissue area. Fresh murine liver tissue were homogenized mechanically and lysed in buffer (Tris-HCL [10mM], NaCl [100mM], EDTA [5mM], Triton-X100 [5%]) containing protease inhibitor (Roche Diagnostics). Lysates (30 µg/ lane) were subjected to 15% SDS-PAGE (Bio-Rad, München, Germany) and blotted on nitrocellulose membranes (Bio-Rad) for Western blot analyses. Northern blot analyses were performed with total RNA isolated from cryopreserved liver tissue using TIRZOL® (Invitrogen).

Human chimeric uPA/SCID mice

Human liver-chimeric UPA/SCID mice were generated by transplanting one million thawed primary human hepatocytes and housed under specific pathogen-free conditions in accordance with protocols approved by the Ethical Committee of the city and state of Hamburg (permission number G12/015) (60). Levels of human chimerism were determined by measuring human serum albumin (HSA) in mouse serum (Human Albumin ELISA kit, Immunology Consultants Lab, Portland, USA) (61). HBV-infected mice received human lymphotoxin-receptor agonizing (CBE11) or isotype control antibodies intra-peritoneally (2mg/kg, body weight) once per week after macrophage depletion using clodronate liposomes (<http://clodronateliposomes.org>). Liver specimens collected at sacrifice were snap-frozen in 2-methylbutane for molecular analyses.

Chimpanzees

Remaining liver biopsies from a study performed on chimpanzees in 2004 were used (26). The study protocol was approved at the Southwest Foundation for Biomedical Research, San Antonio, TX (IACUC 869 PT, approved in 2004). Briefly, animals were treated with 10 million IU of recombinant human interferon-alpha2a and serial liver biopsies were performed at various time points to monitor induction of hepatic genes. Total RNA was extracted from chimpanzee liver biopsies with Trizol reagent (Life Science) and quantified using the NanoDrop spectrophotometer (NanoDrop Technologies). For each reverse transcription reaction, 360 ng of total RNA was converted into cDNA with Transcripter First Strand cDNA Synthesis kit (Roche, Indianapolis, IN, USA) in 20 ul reaction volume. Taqman assay was used to assess the relative

level of the selected transcripts by normalizing to 18s RNA with the delta/delta Ct calculation method. Primers and probes were designed and synthesized by Integration DNA Technologies (Coraville, IA). All qPCR reactions were performed in a 10 µl volume and run on ABI VIIA 7 (Applied Biosystems, Foster City, CA, USA). Each reaction contained 5 µl 2X TaqMan universal PCR master mix (Applied Biosystems,), 0.5 uM of each primer, 0.25 uM of the probe, and cDNA (equivalent to ~16 ng total RNA). The amplification profile was initiated by 10 minute incubation at 95 °C, followed by two-step amplification of 15 seconds at 95 °C and 60 seconds at 60 °C for 40 cycles. For each treatment, results were expressed relative to the expression of the pre-treated chimpanzee liver biopsies as fold-induction. Forward and reverse primers and probes for the selected genes were designed and are listed in Table 1.

GENES	Ref Seq	PROBES(5'-3')	FORWARD PRIMER(5'-3')	REVERSE PRIMER(5'-3')
APOBEC3A	NM_001 270406.1	/56-FAM/TGC AGT AGC /ZEN/ GCA ATC TCG GCT C/3IABkFQ/	AGA TGG AGT CTG GTA CTG TCG	GAG GCA GGA GAG TAG CGT
APOBEC3B	NM_004 900.4	/56-FAM/CCA CAG ATC /ZEN/ AGA AAT CCG ATG GAG CG/3IABkFQ/	GAC AGG GAC AAG CGT ATC TAA G	TCA CTT CAT AGC ACA GCC AAG
APOBEC3C	NM_014 508	/56-FAM/AAC CCG ATG /ZEN/ AAG GCA ATG TAT CCA GG/3IABkFQ/	CAG AAA AGA GTG GGA CAG GG	ACA GCC AAG TTT CGT TCC G
APOBEC3D	M_152426	/56-FAM/CCA CAG ATC /ZEN/ AGA AAT CCG ATG GAG CG/3IABkFQ/	GAC TGG GAC AAG CGT ATC TAA G	TCA CTT CAT AGC ACA GCC AAG
APOBEC3F	NM_145 298	/56-FAM/TGG GAA TAC /ZEN/ACC TGG CCT CGA AAG /3IABkFQ/	CGA AGT GAA AAC AAA GGG TCC	GGA AAC ACT TGT AAG CAG GC
APOBEC3G	NM_021 822.3	/56-FAM/CGG AAT ACA /ZEN/CCT GGC CTC GAA AGA T/3IABkFQ/	CGA AGT GAA AAC AAA GGG TCC	CAT ACT CCT GGT CAC GAT GC
APOBEC3H	NM_001 166003	/56-FAM/ACG GGT CCA /ZEN/AGC TGC AAA CTT CT/3IABkFQ/	CGA CGG CTT GAG AGG ATA AAG	GGC TAT GAG GCA ACT GAC ATG
TBP	NM_003 194	/56-FAM/TGG GAT TAT /ZEN/ATT CGG CGT TTC GGG C/3IABkFQ/	GAG AGT TCT GGG ATT GTA CCG	ATC CTC ATG ATT ACC GCA GC

Analyses of HCV-infected patients treated with PEG-IFN-α

To investigate the effect of IFN-α treatment on APOBEC RNA expression in the human liver we used previously published microarray data of paired liver biopsies from HCV-infected patients undergoing antiviral therapy (62) (Michael T. Dill, Zuzanna Makowska, Gaia Trincucci, Andreas J. Gruber, Julia E. Vogt, Magdalena Filipowicz, Diego Calabrese, Ilona Krol, Daryl T. Lau,

Luigi Terracciano, Erik van Nimwegen, Volker Roth, and Markus H. Heim. Pegylated interferon- α regulates hepatic gene expression by transient activation of the Jak-STAT pathway. Journal of Clinical Investigation, in press). This study included 18 HCV patients who all had a liver biopsy performed during the diagnostic workup and a second liver biopsy collected 4h, 16h, 48h, 96h, or 144h after the first injection of 1.5 ug/kg body weight pegIFNalpha-2b (PegIntronTM, MSD). The patients were subsequently treated with standard combination therapy of pegIFNalpha-2b and ribavirin.

Primers list

Nr.	Oligoname	Sequence (5'->3')
1	ABCA12 fwd	TCTTTTCTTCGCAATGGTCCT
2	ABCA12 rev	GCTGGCATGACTTCTCTATCAA
3	ABCB4 fwd	ATAGCTCACGGATCAGGTCTC
4	ABCB4 rev	GGATTAGCAGCGACAAGGAAA
5	ADAMTS9 fwd	ATTAGAGACCCTGAGCGAATACG
6	ADAMTS9 rev	GAAGTGGACGTTCGTGGGAA
7	ADH1C fwd	GTTGCACCTCTTAAGGCTCAT
8	ADH1C rev	GTTGCCACTAACACATGCT
9	AICDA fwd	GACTTGTTATCTCGCAATAAGA
10	AICDA rev	AGGTCCCAGTCCGAGATGTA
11	AKR1B10 fwd	GTGACACCAGCACGCATTG
12	AKR1B10 rev	GCATTGAAGGGATAGTCTTCAA
13	ALDOB fwd	GGCAGTTCCGAGAAATCCTCT
14	ALDOB rev	CTCCTTGGTCTAACATTGATTCCC
15	ANKH fwd	TCGTGGCCTACCCCTACCTG
16	ANKH rev	TGGATTCTCCCACAAAGCCC
17	ANLN fwd	ATCTGCTGCAACTATTTGCTCC
18	ANLN rev	TCCTGCTAACACTGCTGCTA
19	AOX1 fwd	TGTCGATCCTGAAACAATGCTG
20	AOX1 rev	GGTATGGGTTGTATCGTGA
21	APOBEC3A fwd	GAGAAGGGACAAGCACATGG
22	APOBEC3A rev	TGGATCCATCAAGTGTCTGG
23	APOBEC3B fwd	GACCCTTGGCCTTCGAC
24	APOBEC3B rev	GCACAGCCCCAGGAGAAG
25	APOBEC3C fwd	AGCGCTTCAGAAAAGAGTGG
26	APOBEC3C rev	AAGTTCGTCCGATCGTTG
27	APOBEC3DE fwd	CACATTCTGCGTGGTCTC
28	APOBEC3DE rev	ACCCAAACGTCAGTCGAATC
29	APOBEC3F fwd	CCGTTGGACGCAAAGAT

30	APOBEC3F rev	CCAGGTGATCTGAAACACTT
31	APOBEC3G fwd	CCGAGGACCGAAGGTTAC
32	APOBEC3G rev	TCCAACAGTGCTGAAATTG
33	APOBEC3H fwd	AGCTGTGGCCAGAACGAC
34	APOBEC3H rev	CGGAATGTTGGCTGTT
35	APOBEC4 fwd	TTCTAACACCTGGAATGTGATCC
36	APOBEC4 rev	TTTACTGTCTTAGCTGCAAACC
37	APOC3 fwd	TCAGACCAACTTCAGCCGTG
38	APOC3 rev	GAGCTCGCAGGATGGATAGG
39	ARNT2 fwd	CGTCACCCCTGTTCTGAACC
40	ARNT2 rev	CGGCCTGTCATTGAGTTTCT
41	ASF1B fwd	TCATCCGAGTGGGCTACTACG
42	ASF1B rev	GTTGTTGCCAGTTGATATGGA
43	BID fwd	GTAATCGACCGTGTCCGC
44	BID rev	AACCGTTGTTGACCAACCCTC
45	BIRC3 fwd	TTTCCGTGGCTCTTATTCAAAC
46	BIRC3 rev	GCACAGTGGTAGGAACCTCTCAT
47	C4B fwd	TCCAGGACCTCTCCAGTG
48	C4B rev	CCCCAAAAATGAGCTTGCC
49	C5MTF fwd	GGAAATTAGAACAGGAAATACGA
50	C5MTF rev	AATTGCTTGAGGCGCTTG
51	C6 fwd	TTGATGGGCAATGGGTTCAT
52	C6 rev	ACTTGTCTACTGCTTTGACAG
53	cccDNA 2251 rev	AGCTGAGGCGGTATCTA
54	cccDNA 92 fwd	GCCTATTGATTGAAAGTATGT
55	CCL5 fwd	TCACGCCATTCTCCTG
56	CCL5 rev	CCCCTCACTATCCTACC
57	CCNA2 fwd	GGATGGTAGTTTGAGTCACCAC
58	CCNA2 rev	CACGAGGATAGCTCTCATACTGT
59	CD44 fwd	GCATCGGATTGAGACCTGC
60	CD44 rev	CTGAGACTTGCTGCCCTCTC
61	CD74 fwd	CCGGCTGGACAAACTGACA
62	CD74 rev	GGTGCATCACATGGTCCTCTG
63	CDC20 fwd	GCACAGTTCGCGTTCGAGA
64	CDC20 rev	CTGGATTGCCAGGAGTTGG
65	CKAP2 fwd	GTCCATCAGCATTAAACTCCAGC
66	CKAP2 rev	GTCAGCCTTGGCCAGAAGT
67	CKS2 fwd	CACTACGAGTACGGCATGT
68	CKS2 rev	CACCAAGTCTCCTCCACTCC
69	cMyc-fwd	TCGAGAAGGGCAGGGCTCTCAGAGGCTG
70	cMyc-rev	GGCGATATCGGGTCCACTCCAAGGAGCT
71	CREB3L3 fwd	ATGAATACGGATTAGCTGCTGG

72	CREB3L3 rev	AGGAAGTCGTCAAGAGTCGGG
73	CTSE fwd	AGGCATCCGTCCCTAAGAA
74	CTSE rev	CCTTGGCACTCTGGTCCATTG
75	CX3CL1 fwd	ACCACGGTGTGACGAAATG
76	CX3CL1 rev	TGTTGATAGTGGATGAGCAAAGC
77	CYP2B6 fwd	GCACTCCTCACAGGACTCTG
78	CYP2B6 rev	CCCAGGTGTACCGTGAAGAC
79	CYP3A4 fwd	TTTTTGGATCCATTCTTCTCTCAA
80	CYP3A4 rev	TCCACTCGGTGCTTTGTGT
81	CYP4B1 fwd	CCCACTCTGGTCGGACAG
82	CYP4B1 rev	ATACACATCAGGGGCCTTAGG
83	CYP4F3 fwd	CCCCGAAACGGAATTGGTTCT
84	CYP4F3 rev	TGTGTGTATAGGAGACCTCCTC
85	CYP7A1 fwd	GAGAAGGCAAACGGGTGAAC
86	CYP7A1 rev	GGATTGGCACCAAATTGCAGA
87	DCLK1 fwd	GCTGATTGACCCGAACTCTG
88	DCLK1 rev	AGCCACATACATAACTCTCCT
89	DIO1 fwd	GTCGTGGTAAAGTGCTTCTG
90	DIO1 rev	GTTCCGCTTGACTCTGTCTGG
91	DTL fwd	TCACTGGAATGCCGTCTTGA
92	DTL rev	CTCACCAAGCTTTACGTCCC
93	ECHDC2 fwd	CGACGAUTGAGTGGAACTGAG
94	ECHDC2 rev	CCACAGCGTATTACCAAG
95	EHHADH fwd	AAACTCAGACCCGGTTGAAGA
96	EHHADH rev	TTGCAGAGTCTACGGGATTCT
97	ETNK2 fwd	GCCCCGGCTTTCAAGGTTAAT
98	ETNK2 rev	GGCTGGGTTGATCTCGTT
99	F13B fwd	TCAACCAACCTGTAGGAAAGAAC
100	F13B rev	AGCCAGTAGCACATTGTATTG
101	G6PC fwd	CCCAGGTTCACCAAGTTCCC
102	G6PC rev	GCCGTCAATTGGGCCAGA
103	GAPDH fwd	ACCAACTGCTTAGCCC
104	GAPDH rev	CCACGACGGACACATT
105	GNMT fwd	CTGGGGTGGACTCCATTATGC
106	GNMT rev	GATGACCCACTTGTGAGGC
107	GPC fwd	GTGGCCTCTCAGCCTCG
108	GPC rev	GGCTCTGCAATGGTGGTAGT
109	HAO2 fwd	GCATCACCGGGATGACAA
110	HAO2 rev	GCGATACAAATAGGGGCACTGA
111	HBxin fwd	ATGGCTGCTARGCTGTGCTGCCAA
112	HBxin rev	AAGTGCACACGGTYGGCAGAT
113	HIST13A fwd	ACTGCTCGGAAGTCTACTGGT

114	HIST13A rev	GCGCTGGAAAGGTAGTTACGA
115	HJURP fwd	CCACGCTGACCTACGAGAC
116	HJURP rev	CTCACCGCTTTTGAATCGGC
117	HRG fwd	GACGGGATGGCTACCTTTCC
118	HRG rev	CCGATTCTGCACATCTAAGACT
119	HSD11B1 fwd	AGCAGGAAAGCTCATGGGAG
120	HSD11B1 rev	CCACGTAACTGAGGAAGTTGAC
121	IFNLR2 fwd	ACTGGCTCCCTGGAAGAGAT
122	IFNLR2 rev	TGCAGGGACAGGCCTTTAG
123	IKZF3 fwd	AAAGTGCAGCGGTTTGAATG
124	IKZF3 rev	AGCTGTAAGGGATTCAGGCT
125	ITGA3 fwd	CCCTGGCAGACCTGAACAAT
126	ITGA3 rev	AGACATAGATGGCACCCCCCT
127	ITGAV fwd	ATCTGTGAGGTCGAAACAGGA
128	ITGAV rev	TGGAGCATACTCAACAGTCTTG
129	Kan-L	GGCAGGATCTCCTGTCATCT
130	Kan-R	CATCAGCCATGATGGATACTTC
131	KIF20A fwd	ACCTGATCTGAAGCCCTTGC
132	KIF20A rev	TGGTGCTGGTACCTATCCGA
133	LCN2 fwd	GAAGTGTGACTACTGGATCAGGA
134	LCN2 rev	ACCACTCGGACGAGGTAAC
135	LIPG fwd	GATGGACGATGAGCGGTATCT
136	LIPG rev	CGCATCCGTGAAAGCTGG
137	MBD4 fwd	GGCAACGACTCTAACGAAT
138	MBD4 rev	CCCAAAGCCAGTCATGATATT
139	MKI67 fwd	TTCGCAAGCGCATAACCCA
140	MKI67 rev	AACC GTGTCACAGTGCCAAA
141	MTTP fwd	ACAAGCTCACGTACTCCACTG
142	MTTP rev	TCCTCCATAGTAAGGCCACATC
143	MxA fwd	GGTGGTCCCCAGTAATGTGG
144	MxA rev	CGTCAAGATTCCGATGGTCCT
145	NEIL3 fwd	TCTCCTGTTTGGAAAGTGCA
146	NEIL3 rev	CATTAGCACATCACCTAGCATCC
147	NFE2L3 fwd	TTCAGCCAGGCTATAAGTCAGG
148	NFE2L3 rev	GCTCAGGATTGGTGGTATGAGA
149	NFKB2 fwd	GGGCCGAAAGACCTATCCC
150	NFKB2 rev	CAGCTCCGAGCATTGCTTG
151	NFKBIE fwd	GGTCCACCTGGCAGTGATT
152	NFKBIE rev	ACCATGCCGGTCCTGTAGT
153	NR0B2 fwd	CCCCAAGGAATATGCCTGCC
154	NR0B2 rev	TAGGGCGAAAGAAGAGGTCCC
155	NR1D1 fwd	TGGACTCCAACAACACACAG

156	NR1D1 rev	GATGGTGGGAAGTAGGTGGG
157	NR1I3 fwd	TGGGCACCATGTTAACAGT
158	NR1I3 rev	GGGCAGGTCTTAGTAAACTTG
159	OAS1 fwd	TGTCCAAGGTGGTAAAGGGTG
160	OAS1 rev	CCGGCGATTAACTGATCCTG
161	PFKFB1 fwd	AGAAGGGGCTCATCCATACCC
162	PFKFB1 rev	CTCTCGTCGATACTGGCCTAA
163	pgRNA 383 fwd	CTCCTCCAGCTTATAGACC
164	pgRNA 705 rev	GTGAGTGGGCCTACAAA
165	PIPOX fwd	GCAGAGGGTAGAACACCAGTG
166	PIPOX rev	CCTAGCTGTCGAATTGCATCC
167	PKR fwd	GCCGCTAAACTGCATATCTCA
168	PKR rev	TCACACGTAGTAGCAAAAGAACCC
169	PLA2G2A fwd	ATGAAGACCCCTCCTACTGTTGG
170	PLA2G2A rev	GCTTCCTTCCTGTCGTCAACT
171	PLK1 fwd	CGAGGACAACGACTCGTGT
172	PLK1 rev	ACAATTGCCGTAGGTAGTATCG
173	PNPLA3 fwd	GGCATCTCTCTTACCAAGAGTGT
174	PNPLA3 rev	GGCATCCACGACTCGTCTTT
175	POR fwd	GGTGGCCGAAGAAGTATCTCT
176	POR rev	AACCAGTAGGTTAGGAGACCC
177	Prnp fwd	TGCTGGGAAGTGCATGAG
178	Prnp rev	CGGTGCATGTTTACGATAGTA
179	PSMB9 fwd	GGTTCTGATTCCCGAGTGTCT
180	PSMB9 rev	CAGCCAAAACAAGTGGAGGTT
181	p53-fwd	GAGTGCAGTGGCACGATT
182	p53-rev	GAATCGCTTCAGCTCAGGA
183	RAC2fwd	CAACGCCCTTCCGGAGAG
184	RAC2rev	TCCGTCTGGATAGGAGAGC
185	rcDNA1745 fwd	GTTGCCGTTGTCCTCTAATT
186	rcDNA1844 rev	GGAGGGATACATAGAGGTTCTGA
187	RCN1 fwd	TGGATCAAACGGGTGCAGAA
188	RCN1 rev	TGCGGGTTCTAGGTAGT
189	RDH5 fwd	TGGGTGGAGATGCACGTTAAG
190	RDH5 rev	GTGTGGTCCGATGATACCAAG
191	RRM2 fwd	ATTGGGCCTTGCATGGATAG
192	RRM2 rev	GAGTCCTGGCATAAGACCTCT
193	SCP2 fwd	TACCCCTGACTTGGCAGAAGAA
194	SCP2 rev	CTGGCGGGCCATAAACAAA
195	SERPINA7 fwd	TCACCATTCCTGTATCTGGTCT
196	SERPINA7 rev	GTGGCATTGGTTGGATGAA
197	SERPIN D1 fwd	GGACGACGACTATCTGGACCT

198	SERPIND1 rev	CATCAGAGTCTGTCGGGGAAA
199	SERPING1 fwd	CCAAGATGCTATTGTTGAACCC
200	SERPING1 rev	TGGTGGCTGAATTGGTTGTTG
201	SLC17A4 fwd	ATGTCTACCGGACCAGATGTC
202	SLC17A4 rev	GCCCATGTCGGACTGAACAAA
203	SLC22A1 fwd	TGCCCTCATTTGTTGCG
204	SLC22A1 rev	TTTCTCCCAGGTTCTCGGC
205	SLC9A9 fwd	CAGCACAAACATCAATCCTCATCA
206	SLC9A9 rev	CAGCATGTATCATAGCCTTCACA
207	SMUG1 fwd	CTGCAGTGCCTGTATGTG
208	SMUG1 rev	GCAGGCTCATGGATGGAC
209	SOD2 fwd	GGAAGCCATCAAACGTGACTT
210	SOD2 rev	CCCGTTCCATTGAAACCAAGC
211	SPP1 fwd	CTCCATTGACTCGAACGACTC
212	SPP1 rev	CAGGTCTCGCAAACCTTAGAT
213	Src-fw	TGTGTGTGAGAAAACACAAAAT
214	Src-rev	TCAGTCCTCCCAGGATGTTC
215	SULT2A1 fwd	AGCGATCACCTGGTAGAG
216	SULT2A1 rev	TGGGAGGAGAATAAACGTGGAC
217	TAP1fwd	CGCCTCACTGACTGGATTCTA
218	TAP1rev	TCTGTTGGAAAAACTCCGTCTC
219	TAT fwd	CTGGACTCGGGCAAATATAATGG
220	TAT rev	GTCCTTAGCTTAGGGGTGC
221	TDG fwd	GAACCTTGTGGCTCTCTCA
222	TDG rev	GTCATCCACTGCCATTAGG
223	TF fwd	GTGTGCAGTGTGGAGCAT
224	Tf rev	CATCGGATGGAATGACGCTT
225	TMEM123 fwd	GGCGGCATCTGCAAACATAG
226	TMEM123 rev	TTCATGGTGGTGACCGTTGT
227	TMPRSS2 fwd	CAAGTGCTCAACTCTGGGAT
228	TMPRSS2 rev	AACACACCGATTCTCGTCCTC
229	TNFAIP3 fwd	TCAACTGGTGTGAGAAGTCC
230	TNFAIP3 rev	CAAGTCTGTGCTCTGAACGC
231	TNFRSF9 fwd	TCCACCAAGCAATGCAGAGTG
232	TNFRSF9 rev	CCAAAGCAACAGTCTTACAACC
233	TOP2A fwd	ACCATTGCAGCCTGTAAATGA
234	TOP2A rev	GGGCGGAGCAAATATGTTCC
235	TPX2 fwd	ACTTCCGCACAGATGAGCG
236	TPX2 rev	GGATGCTTCGTAGTCAGATGT
237	TRAF1 fwd	CGCCGAGATGGAGTCATCAG
238	TRAF1 rev	TGGAAATCCCCAGGGTGTG
239	TYMS fwd	CTGCTGACAACCAAACGTGTG

240	TYMS rev	GCATCCCAGATTTCACTCCCTT
241	UBD fwd	CCGTTCCGAGGAATGGGATT
242	UBD rev	GCCATAAGATGAGAGGCTCTCC
244	UGT2B10 fwd	GAAATGGACTACAGTTCTGCTGA
245	UGT2B10 rev	GTGGATGAGTCGTTGGGATCA
246	UGT2B4 fwd	CAAATGTTGAGTCGTTGGAGGA
247	UGT2B4 rev	CTGACGTGTTACTGACCATCG
248	V-MYC fwd	GGCTCCTGGCAAAAGGTCA
249	V-MYC rev	CTGCGTAGTTGTGCTGATGT
250	VTN fwd	CGGGGATGTGTTCACTATGCC
251	VTN rev	GTGTCTGCTCAGGATTCCCTT

qRT-PCR primer sequences were either taken from the Harvard primer bank or newly established by using NCBI primer blast. All primer pairs were designed to span at least one intron/exon border to exclude genomic DNA amplification and checked for sequence specificity to avoid unspecific by-products. Correct amplicon size was checked by agarose gel-electrophoreses and melting curve analyses were done for each qPCR run.

References

1. U. Protzer, M. K. Maini, P. A. Knolle, Living in the liver: hepatic infections. *Nat. Rev. Immunol.* **12**, 201–213 (2012). [doi:10.1038/nri3169](https://doi.org/10.1038/nri3169) [Medline](#)
2. F. Zoulim, Hepatitis B virus resistance to antiviral drugs: where are we going? *Liver Int.* **31**, (Suppl 1), 111–116 (2011). [doi:10.1111/j.1478-3231.2010.02399.x](https://doi.org/10.1111/j.1478-3231.2010.02399.x) [Medline](#)
3. K. Wursthorn, M. Lutgehetmann, M. Dandri, T. Volz, P. Buggisch, B. Zollner, T. Longerich, P. Schirmacher, F. Metzler, M. Zankel, C. Fischer, G. Currie, C. Brosgart, J. Petersen, Peginterferon alpha-2b plus adefovir induce strong cccDNA decline and HBsAg reduction in patients with chronic hepatitis B. *Hepatology* **44**, 675–684 (2006). [doi:10.1002/hep.21282](https://doi.org/10.1002/hep.21282) [Medline](#)
4. L. G. Guidotti, K. Ando, M. V. Hobbs, T. Ishikawa, L. Runkel, R. D. Schreiber, F. V. Chisari, Cytotoxic T lymphocytes inhibit hepatitis B virus gene expression by a noncytolytic mechanism in transgenic mice. *Proc. Natl. Acad. Sci. U.S.A.* **91**, 3764–3768 (1994). [doi:10.1073/pnas.91.9.3764](https://doi.org/10.1073/pnas.91.9.3764) [Medline](#)
5. L. G. Guidotti, R. Rochford, J. Chung, M. Shapiro, R. Purcell, F. V. Chisari, Viral clearance without destruction of infected cells during acute HBV infection. *Science* **284**, 825–829 (1999). [doi:10.1126/science.284.5415.825](https://doi.org/10.1126/science.284.5415.825) [Medline](#)
6. S. F. Wieland, H. C. Spangenberg, R. Thimme, R. H. Purcell, F. V. Chisari, Expansion and contraction of the hepatitis B virus transcriptional template in infected chimpanzees. *Proc. Natl. Acad. Sci. U.S.A.* **101**, 2129–2134 (2004). [doi:10.1073/pnas.0308478100](https://doi.org/10.1073/pnas.0308478100) [Medline](#)
7. H. McClary, R. Koch, F. V. Chisari, L. G. Guidotti, Relative sensitivity of hepatitis B virus and other hepatotropic viruses to the antiviral effects of cytokines. *J. Virol.* **74**, 2255–2264 (2000). [doi:10.1128/JVI.74.5.2255-2264.2000](https://doi.org/10.1128/JVI.74.5.2255-2264.2000) [Medline](#)
8. L. Belloni, L. Allweiss, F. Guerrieri, N. Pediconi, T. Volz, T. Pollicino, J. Petersen, G. Raimondo, M. Dandri, M. Leviero, IFN- α inhibits HBV transcription and replication in cell culture and in humanized mice by targeting the epigenetic regulation of the nuclear cccDNA minichromosome. *J. Clin. Invest.* **122**, 529–537 (2012). [doi:10.1172/JCI58847](https://doi.org/10.1172/JCI58847) [Medline](#)

9. A. Rang, S. Günther, H. Will, Effect of interferon alpha on hepatitis B virus replication and gene expression in transiently transfected human hepatoma cells. *J. Hepatol.* **31**, 791–799 (1999). [doi:10.1016/S0168-8278\(99\)80279-7](https://doi.org/10.1016/S0168-8278(99)80279-7) [Medline](#)
10. V. Pasquetto, S. F. Wieland, S. L. Uprichard, M. Tripodi, F. V. Chisari, Cytokine-sensitive replication of hepatitis B virus in immortalized mouse hepatocyte cultures. *J. Virol.* **76**, 5646–5653 (2002). [doi:10.1128/JVI.76.11.5646-5653.2002](https://doi.org/10.1128/JVI.76.11.5646-5653.2002) [Medline](#)
11. S. F. Wieland, L. G. Guidotti, F. V. Chisari, Intrahepatic induction of alpha/beta interferon eliminates viral RNA-containing capsids in hepatitis B virus transgenic mice. *J. Virol.* **74**, 4165–4173 (2000). [doi:10.1128/JVI.74.9.4165-4173.2000](https://doi.org/10.1128/JVI.74.9.4165-4173.2000) [Medline](#)
12. S. L. Uprichard, S. F. Wieland, A. Althage, F. V. Chisari, Transcriptional and posttranscriptional control of hepatitis B virus gene expression. *Proc. Natl. Acad. Sci. U.S.A.* **100**, 1310–1315 (2003). [doi:10.1073/pnas.252773599](https://doi.org/10.1073/pnas.252773599) [Medline](#)
13. P. Gripon, C. Diot, N. Thézé, I. Fourel, O. Loreal, C. Brechot, C. Guguen-Guillouzo, Hepatitis B virus infection of adult human hepatocytes cultured in the presence of dimethyl sulfoxide. *J. Virol.* **62**, 4136–4143 (1988). [Medline](#)
14. P. Gripon, S. Rumin, S. Urban, J. Le Seyec, D. Glaise, I. Cannie, C. Guyomard, J. Lucas, C. Trepo, C. Guguen-Guillouzo, Infection of a human hepatoma cell line by hepatitis B virus. *Proc. Natl. Acad. Sci. U.S.A.* **99**, 15655–15660 (2002). [doi:10.1073/pnas.232137699](https://doi.org/10.1073/pnas.232137699) [Medline](#)
15. M. J. Wolf, G. M. Seleznik, N. Zeller, M. Heikenwalder, The unexpected role of lymphotoxin beta receptor signaling in carcinogenesis: from lymphoid tissue formation to liver and prostate cancer development. *Oncogene* **29**, 5006–5018 (2010). [doi:10.1038/onc.2010.260](https://doi.org/10.1038/onc.2010.260) [Medline](#)
16. J. Haybaeck, N. Zeller, M. J. Wolf, A. Weber, U. Wagner, M. O. Kurrer, J. Bremer, G. Iezzi, R. Graf, P. A. Clavien, R. Thimme, H. Blum, S. A. Nedospasov, K. Zatloukal, M. Ramzan, S. Ciesek, T. Pietschmann, P. N. Marche, M. Karin, M. Kopf, J. L. Browning, A. Aguzzi, M. Heikenwalder, A lymphotoxin-driven pathway to hepatocellular carcinoma. *Cancer Cell* **16**, 295–308 (2009). [doi:10.1016/j.ccr.2009.08.021](https://doi.org/10.1016/j.ccr.2009.08.021) [Medline](#)

17. S. Jost, P. Turelli, B. Mangeat, U. Protzer, D. Trono, Induction of antiviral cytidine deaminases does not explain the inhibition of hepatitis B virus replication by interferons. *J. Virol.* **81**, 10588–10596 (2007). [doi:10.1128/JVI.02489-06](https://doi.org/10.1128/JVI.02489-06) [Medline](#)
18. M. Lukashev, D. LePage, C. Wilson, V. Bailly, E. Garber, A. Lukashin, A. Ngam-ek, W. Zeng, N. Allaire, S. Perrin, X. Xu, K. Szeliga, K. Wortham, R. Kelly, C. Bottiglio, J. Ding, L. Griffith, G. Heaney, E. Silverio, W. Yang, M. Jarpe, S. Fawell, M. Reff, A. Carmillo, K. Miatkowski, J. Amatucci, T. Crowell, H. Prentice, W. Meier, S. M. Violette, F. Mackay, D. Yang, R. Hoffman, J. L. Browning, Targeting the lymphotoxin-beta receptor with agonist antibodies as a potential cancer therapy. *Cancer Res.* **66**, 9617–9624 (2006). [doi:10.1158/0008-5472.CAN-06-0217](https://doi.org/10.1158/0008-5472.CAN-06-0217) [Medline](#)
19. X. Hu, M. A. Zimmerman, K. Bardhan, D. Yang, J. L. Waller, G. B. Liles, J. R. Lee, R. Pollock, D. Lev, C. F. Ware, E. Garber, V. Bailly, J. L. Browning, K. Liu, Lymphotoxin β receptor mediates caspase-dependent tumor cell apoptosis in vitro and tumor suppression in vivo despite induction of NF- κ B activation. *Carcinogenesis* **34**, 1105–1114 (2013). [doi:10.1093/carcin/bgt014](https://doi.org/10.1093/carcin/bgt014) [Medline](#)
20. E. Dejardin, N. M. Droin, M. Delhase, E. Haas, Y. Cao, C. Makris, Z. W. Li, M. Karin, C. F. Ware, D. R. Green, The lymphotoxin-beta receptor induces different patterns of gene expression via two NF-kappaB pathways. *Immunity* **17**, 525–535 (2002). [doi:10.1016/S1074-7613\(02\)00423-5](https://doi.org/10.1016/S1074-7613(02)00423-5) [Medline](#)
21. R. Suspène, D. Guétard, M. Henry, P. Sommer, S. Wain-Hobson, J. P. Vartanian, Extensive editing of both hepatitis B virus DNA strands by APOBEC3 cytidine deaminases in vitro and in vivo. *Proc. Natl. Acad. Sci. U.S.A.* **102**, 8321–8326 (2005). [doi:10.1073/pnas.0408223102](https://doi.org/10.1073/pnas.0408223102) [Medline](#)
22. J. I. Friedman, J. T. Stivers, Detection of damaged DNA bases by DNA glycosylase enzymes. *Biochemistry* **49**, 4957–4967 (2010). [doi:10.1021/bi100593a](https://doi.org/10.1021/bi100593a) [Medline](#)
23. M. D. Stenglein, M. B. Burns, M. Li, J. Lengyel, R. S. Harris, APOBEC3 proteins mediate the clearance of foreign DNA from human cells. *Nat. Struct. Mol. Biol.* **17**, 222–229 (2010). [doi:10.1038/nsmb.1744](https://doi.org/10.1038/nsmb.1744) [Medline](#)

24. M. Bonvin, F. Achermann, I. Greeve, D. Stroka, A. Keogh, D. Inderbitzin, D. Candinas, P. Sommer, S. Wain-Hobson, J. P. Vartanian, J. Greeve, Interferon-inducible expression of APOBEC3 editing enzymes in human hepatocytes and inhibition of hepatitis B virus replication. *Hepatology* **43**, 1364–1374 (2006). [doi:10.1002/hep.21187](https://doi.org/10.1002/hep.21187) [Medline](#)
25. T. Zhang, J. Cai, J. Chang, D. Yu, C. Wu, T. Yan, K. Zhai, X. Bi, H. Zhao, J. Xu, W. Tan, C. Qu, D. Lin, Evidence of associations of APOBEC3B gene deletion with susceptibility to persistent HBV infection and hepatocellular carcinoma. *Hum. Mol. Genet.* **22**, 1262–1269 (2013). [doi:10.1093/hmg/ddt513](https://doi.org/10.1093/hmg/ddt513) [Medline](#)
26. Y. Huang, J. J. Feld, R. K. Sapp, S. Nanda, J. H. Lin, L. M. Blatt, M. W. Fried, K. Murthy, T. J. Liang, Defective hepatic response to interferon and activation of suppressor of cytokine signaling 3 in chronic hepatitis C. *Gastroenterology* **132**, 733–744 (2007). [doi:10.1053/j.gastro.2006.11.045](https://doi.org/10.1053/j.gastro.2006.11.045) [Medline](#)
27. B. P. Doehle, A. Schäfer, B. R. Cullen, Human APOBEC3B is a potent inhibitor of HIV-1 infectivity and is resistant to HIV-1 Vif. *Virology* **339**, 281–288 (2005). [doi:10.1016/j.virol.2005.06.005](https://doi.org/10.1016/j.virol.2005.06.005) [Medline](#)
28. G. Berger, J. Turpin, S. Cordeil, K. Tartour, X. N. Nguyen, R. Mahieux, A. Cimarelli, Functional analysis of the relationship between Vpx and the restriction factor SAMHD1. *J. Biol. Chem.* **287**, 41210–41217 (2012). [doi:10.1074/jbc.M112.403816](https://doi.org/10.1074/jbc.M112.403816) [Medline](#)
29. H. Muckenfuss, M. Hamdorf, U. Held, M. Perkovic, J. Löwer, K. Cichutek, E. Flory, G. G. Schumann, C. Münk, APOBEC3 proteins inhibit human LINE-1 retrotransposition. *J. Biol. Chem.* **281**, 22161–22172 (2006). [doi:10.1074/jbc.M601716200](https://doi.org/10.1074/jbc.M601716200) [Medline](#)
30. C. Münk, A. Willemse, I. G. Bravo, An ancient history of gene duplications, fusions and losses in the evolution of APOBEC3 mutators in mammals. *BMC Evol. Biol.* **12**, 71 (2012). [doi:10.1186/1471-2148-12-71](https://doi.org/10.1186/1471-2148-12-71) [Medline](#)
31. M. A. Carpenter, M. Li, A. Rathore, L. Lackey, E. K. Law, A. M. Land, B. Leonard, S. M. Shandilya, M. F. Bohn, C. A. Schiffer, W. L. Brown, R. S. Harris, Methylcytosine and normal cytosine deamination by the foreign DNA restriction enzyme APOBEC3A. *J. Biol. Chem.* **287**, 34801–34808 (2012). [doi:10.1074/jbc.M112.385161](https://doi.org/10.1074/jbc.M112.385161) [Medline](#)


32. P. Turelli, B. Mangeat, S. Jost, S. Vianin, D. Trono, Inhibition of hepatitis B virus replication by APOBEC3G. *Science* **303**, 1829 (2004). [doi:10.1126/science.1092066](https://doi.org/10.1126/science.1092066) [Medline](#)
33. C. T. Bock, S. Schwinn, S. Locarnini, J. Fyfe, M. P. Manns, C. Trautwein, H. Zentgraf, Structural organization of the hepatitis B virus minichromosome. *J. Mol. Biol.* **307**, 183–196 (2001). [doi:10.1006/jmbi.2000.4481](https://doi.org/10.1006/jmbi.2000.4481) [Medline](#)
34. M. M. Aynaud, R. Suspène, P. O. Vidalain, B. Mussil, D. Guétard, F. Tangy, S. Wain-Hobson, J. P. Vartanian, Human Tribbles 3 protects nuclear DNA from cytidine deamination by APOBEC3A. *J. Biol. Chem.* **287**, 39182–39192 (2012). [doi:10.1074/jbc.M112.372722](https://doi.org/10.1074/jbc.M112.372722) [Medline](#)
35. Y. Guo, W. Kang, X. Lei, Y. Li, A. Xiang, Y. Liu, J. Zhao, J. Zhang, Z. Yan, Hepatitis B viral core protein disrupts human host gene expression by binding to promoter regions. *BMC Genomics* **13**, 563 (2012). [doi:10.1186/1471-2164-13-563](https://doi.org/10.1186/1471-2164-13-563) [Medline](#)
36. H. C. Smith, R. P. Bennett, A. Kizilyer, W. M. McDougall, K. M. Prohaska, Functions and regulation of the APOBEC family of proteins. *Semin. Cell Dev. Biol.* **23**, 258–268 (2012). [doi:10.1016/j.semcdb.2011.10.004](https://doi.org/10.1016/j.semcdb.2011.10.004) [Medline](#)
37. T. H. Lee, S. J. Elledge, J. S. Butel, Hepatitis B virus X protein interacts with a probable cellular DNA repair protein. *J. Virol.* **69**, 1107–1114 (1995). [Medline](#)
38. K. Kitamura, Z. Wang, S. Chowdhury, M. Simadu, M. Koura, M. Muramatsu, Uracil DNA glycosylase counteracts APOBEC3G-induced hypermutation of hepatitis B viral genomes: excision repair of covalently closed circular DNA. *PLoS Pathog.* **9**, e1003361 (2013). [doi:10.1371/journal.ppat.1003361](https://doi.org/10.1371/journal.ppat.1003361) [Medline](#)
39. S. Landry, I. Narvaiza, D. C. Linfesty, M. D. Weitzman, APOBEC3A can activate the DNA damage response and cause cell-cycle arrest. *EMBO Rep.* **12**, 444–450 (2011). [doi:10.1038/embor.2011.46](https://doi.org/10.1038/embor.2011.46) [Medline](#)
40. M. B. Burns, L. Lackey, M. A. Carpenter, A. Rathore, A. M. Land, B. Leonard, E. W. Refsland, D. Kotandeniya, N. Tretyakova, J. B. Nikas, D. Yee, N. A. Temiz, D. E. Donohue, R. M. McDougle, W. L. Brown, E. K. Law, R. S. Harris, APOBEC3B is an enzymatic source of mutation in breast cancer. *Nature* **494**, 366–370 (2013). [doi:10.1038/nature11881](https://doi.org/10.1038/nature11881) [Medline](#)

41. K. Krebs, N. Böttiger, L. R. Huang, M. Chmielewski, S. Arzberger, G. Gasteiger, C. Jäger, E. Schmitt, F. Bohne, M. Aichler, W. Uckert, H. Abken, M. Heikenwalder, P. Knolle, U. Protzer, T cells expressing a chimeric antigen receptor that binds hepatitis B virus envelope proteins control virus replication in mice. *Gastroenterology* **145**, 456–465 (2013). [doi:10.1053/j.gastro.2013.04.047](https://doi.org/10.1053/j.gastro.2013.04.047) [Medline](#)
42. J. Lucifora, S. Arzberger, D. Durantel, L. Belloni, M. Strubin, M. Levrero, F. Zoulim, O. Hantz, U. Protzer, Hepatitis B virus X protein is essential to initiate and maintain virus replication after infection. *J. Hepatol.* **55**, 996–1003 (2011).
[doi:10.1016/j.jhep.2011.02.015](https://doi.org/10.1016/j.jhep.2011.02.015) [Medline](#)
43. H. Schulze-Bergkamen, A. Untergasser, A. Dax, H. Vogel, P. Büchler, E. Klar, T. Lehnert, H. Friess, M. W. Büchler, M. Kirschfink, W. Stremmel, P. H. Krammer, M. Müller, U. Protzer, Primary human hepatocytes—a valuable tool for investigation of apoptosis and hepatitis B virus infection. *J. Hepatol.* **38**, 736–744 (2003). [doi:10.1016/S0168-8278\(03\)00120-X](https://doi.org/10.1016/S0168-8278(03)00120-X) [Medline](#)
44. S. M. Lee, C. Schelcher, M. Demmel, M. Hauner, W. E. Thasler, Isolation of human hepatocytes by a two-step collagenase perfusion procedure. *J. Vis. Exp.* (79): (2013). [Medline](#)
45. W. E. Thasler, T. S. Weiss, K. Schillhorn, P. T. Stoll, B. Irrgang, K. W. Jauch, Charitable State-Controlled Foundation Human Tissue and Cell Research: Ethic and Legal Aspects in the Supply of Surgically Removed Human Tissue For Research in the Academic and Commercial Sector in Germany. *Cell Tissue Bank.* **4**, 49–56 (2003).
[doi:10.1023/A:1026392429112](https://doi.org/10.1023/A:1026392429112) [Medline](#)
46. O. Hantz, R. Parent, D. Durantel, P. Gripon, C. Guguen-Guillouzo, F. Zoulim, Persistence of the hepatitis B virus covalently closed circular DNA in HepaRG human hepatocyte-like cells. *J. Gen. Virol.* **90**, 127–135 (2009). [doi:10.1099/vir.0.004861-0](https://doi.org/10.1099/vir.0.004861-0) [Medline](#)
47. M. Quasdorff, M. Hösel, M. Odenthal, U. Zedler, F. Bohne, P. Gripon, H. P. Dienes, U. Drebber, D. Stippel, T. Goeser, U. Protzer, A concerted action of HNF4alpha and HNF1alpha links hepatitis B virus replication to hepatocyte differentiation. *Cell. Microbiol.* **10**, 1478–1490 (2008). [doi:10.1111/j.1462-5822.2008.01141.x](https://doi.org/10.1111/j.1462-5822.2008.01141.x) [Medline](#)


48. U. Protzer, S. Seyfried, M. Quasdorff, G. Sass, M. Svorcova, D. Webb, F. Bohne, M. Hösel, P. Schirmacher, G. Tiegs, Antiviral activity and hepatoprotection by heme oxygenase-1 in hepatitis B virus infection. *Gastroenterology* **133**, 1156–1165 (2007). [doi:10.1053/j.gastro.2007.07.021](https://doi.org/10.1053/j.gastro.2007.07.021) [Medline](#)
49. A. Untergasser, U. Zedler, A. Langenkamp, M. Hösel, M. Quasdorff, K. Esser, H. P. Dienes, B. Tappertzhofen, W. Kolanus, U. Protzer, Dendritic cells take up viral antigens but do not support the early steps of hepatitis B virus infection. *Hepatology* **43**, 539–547 (2006). [doi:10.1002/hep.21048](https://doi.org/10.1002/hep.21048) [Medline](#)
50. J. Summers, P. M. Smith, A. L. Horwich, Hepadnavirus envelope proteins regulate covalently closed circular DNA amplification. *J. Virol.* **64**, 2819–2824 (1990). [Medline](#)
51. W. Gao, J. Hu, Formation of hepatitis B virus covalently closed circular DNA: removal of genome-linked protein. *J. Virol.* **81**, 6164–6174 (2007). [doi:10.1128/JVI.02721-06](https://doi.org/10.1128/JVI.02721-06) [Medline](#)
52. G. K. Smyth, Linear models and empirical bayes methods for assessing differential expression in microarray experiments. *Stat. Appl. Genet. Mol. Biol.* **3**, Article3 (2004).
53. C. Banning, J. Votteler, D. Hoffmann, H. Koppensteiner, M. Warmer, R. Reimer, F. Kirchhoff, U. Schubert, J. Hauber, M. Schindler, A flow cytometry-based FRET assay to identify and analyse protein-protein interactions in living cells. *PLoS ONE* **5**, e9344 (2010). [doi:10.1371/journal.pone.0009344](https://doi.org/10.1371/journal.pone.0009344) [Medline](#)
54. K. Arnold, L. Bordoli, J. Kopp, T. Schwede, The SWISS-MODEL workspace: a web-based environment for protein structure homology modelling. *Bioinformatics* **22**, 195–201 (2006). [doi:10.1093/bioinformatics/bti770](https://doi.org/10.1093/bioinformatics/bti770) [Medline](#)
55. S. A. Wynne, R. A. Crowther, A. G. Leslie, The crystal structure of the human hepatitis B virus capsid. *Mol. Cell* **3**, 771–780 (1999). [doi:10.1016/S1097-2765\(01\)80009-5](https://doi.org/10.1016/S1097-2765(01)80009-5) [Medline](#)
56. I. J. Byeon *et al.*, NMR structure of human restriction factor APOBEC3A reveals substrate binding and enzyme specificity. *Nat. Commun.* **4**, 1890 (2013).
57. A. W. Ghoorah, M. D. Devignes, M. Smail-Tabbone, D. W. Ritchie, Protein docking using case-based reasoning. *Proteins* **81**, 2150–2158 (2013). [doi:10.1002/prot.24433](https://doi.org/10.1002/prot.24433) [Medline](#)

58. R. A. Sayle, E. J. Milner-White, RASMOL: biomolecular graphics for all. *Trends Biochem. Sci.* **20**, 374–376 (1995). [doi:10.1016/S0968-0004\(00\)89080-5](https://doi.org/10.1016/S0968-0004(00)89080-5) [Medline](#)
59. L. G. Guidotti, B. Matzke, H. Schaller, F. V. Chisari, High-level hepatitis B virus replication in transgenic mice. *J. Virol.* **69**, 6158–6169 (1995). [Medline](#)
60. M. Dandri, M. R. Burda, E. Török, J. M. Pollok, A. Iwanska, G. Sommer, X. Rogiers, C. E. Rogler, S. Gupta, H. Will, H. Greten, J. Petersen, Repopulation of mouse liver with human hepatocytes and in vivo infection with hepatitis B virus. *Hepatology* **33**, 981–988 (2001). [doi:10.1053/jhep.2001.23314](https://doi.org/10.1053/jhep.2001.23314) [Medline](#)
61. M. Lütgehetmann, T. Bornscheuer, T. Volz, L. Allweiss, J. H. Bockmann, J. M. Pollok, A. W. Lohse, J. Petersen, M. Dandri, Hepatitis B virus limits response of human hepatocytes to interferon- α in chimeric mice. *Gastroenterology* **140**, 2074–2083, e1–e2 (2011). [doi:10.1053/j.gastro.2011.02.057](https://doi.org/10.1053/j.gastro.2011.02.057) [Medline](#)
62. M. Sarasin-Filipowicz, E. J. Oakeley, F. H. Duong, V. Christen, L. Terracciano, W. Filipowicz, M. H. Heim, Interferon signaling and treatment outcome in chronic hepatitis C. *Proc. Natl. Acad. Sci. U.S.A.* **105**, 7034–7039 (2008). [doi:10.1073/pnas.0707882105](https://doi.org/10.1073/pnas.0707882105) [Medline](#)


A




B



C



D



E




Figure S1. (A) The indicated cells were treated with IFN- α for 6 hours and levels of Mx1 and OAS1 were assessed by qRT-PCR. (B, C) dHepaRG cells were infected with HBV and treated 10 days later with IFN- α or Lamivudine (LAM). (B) Viability of cells was assessed by XTT test. (C) Levels of HBeAg, intracellular HBV-DNA, HBV-RNA or HBV-DNA in the supernatants were analyzed and compared to untreated cells (mock). Values given are the mean +/- standard deviation of biological triplicates and were analyzed by t-test. ** p<0.01 and *** p<0.001. (D) HBV cccDNA levels were analyzed by Southern blot 10 days post-treatment. Supercoiled cccDNA bands were identified by their expected size and ability to be linearized upon EcoRI digestion (3.2 kb). (E) Dose response curve of viral markers after IFN- α treatment for 7 days.









A**B****C****E****D****F****G****H**

Figure S2. HBV-infected dHepaRG cells were treated with BS1. After 10 days, we analyzed **(A)** expression of LT β R and TNFR1 by Western blot and **(B)** by FACS analyses as well as **(C)** p100/p52 processing and **(D)** phosphorylation of RelA (pRelA) by Western blot. **(E)** Nuclear translocation of RelB and of **(F)** RelA by immunohistochemistry (white arrow heads indicate nuclear translocation in hepatocytes). **(G)** Expression analysis of indicated genes upon BS1 treatment by qRT-PCR. **(H)** Assessment of cell viability under different concentration of BS1 by XTT tests.





A**B****C****D**

Figure S3. dHepaRG cells were infected with HBV and treated with LT β R-agonist BS1. **(A)** Dose response curve of viral markers upon BS1 treatment. **(B)** cccDNA signal detected by Southern blot analysis was quantified using a phosphorimager and compared to quantification of cccDNA of the same extract by qPCR. **(C, D)** HBV-infected dHepaRG cells were treated with BS1 as indicated in the schemes. cccDNA was quantified by qPCR relative to untreated cells (mock). Values are the mean +/- standard deviation of different replicates and data were analyzed by t-test. ** $p < 0.01$ and *** $p < 0.001$.




Figure S4. dHepaRG cells or primary human hepatocytes (PHH) were infected with HBV and treated with BS1 for 11 days and assayed for apoptosis induction. As positive control, cells were treated 12h before the end of the experiment with 1 μ g/mL of Staurosporine (Stauro). ALT (**A**) and HBeAg levels (**B, right panel**) were quantified in the supernatant. Cleaved caspase 3 activity was analyzed (**B, left panel**).





Figure S5. To exclude induction of IFN responses, HepaRG cells expressing a dominant negative version of the IFN regulated factor 3 (IRF3) were generated. (**A**) dHepaRG and dHepaRG-IRF3DN cells were stimulated with poly-IC during 8h and expression of IFN- β was analyzed by qRT-PCR (relative to GAPDH). (**B**) dHepaRG or dHepaRG-IRF3DN cells were infected with HBV and treated with BS1 24h later and for 10 days. cccDNA content was analyzed and expressed in % of untreated cells (mock). Values are given as mean +/- standard deviation of three replicates and data were analyzed by t-test. ** p<0.01 and *** p<0.001.










A**B****C****D**


Figure S7. HBV-infected dHepaRG cells were treated with **(A)** IFN- α or **(B)** LT β R-agonist BS1, for 10 days. 3D-PCRs analyses were performed on cccDNA. **(A, B)** 3D-PCR products obtained at indicated temperatures were cloned, sequenced and numbers of G/A mutations were evaluated. **(C)** 3D-PCR of cccDNA after 10 days of treatment with IFN- α or nucleoside analogues lamivudine (LAM) or entecavir (ETV). **(D)** Treatment with IFN- α alone was compared to combined IFN- α +ETV treatment.

A



B

Mock
IFN
BS1



C

Gene Name	Amplicon Length	Genomic Position	% GC	Average coverage	% overall Mutation Rate (Mock/IFN/BS1)	G-A Mutation Rate (Mock/IFN/BS1)	C-T Mutation Rate (Mock/IFN/BS1)
B2M	2157	Chr. 15: 15798407 - 15800563	40,10	85 543	0.31/0.31/0.31	0.050/0.045/0.045	0.055/0.050/0.050
TBP	2249	Chr. 6: 75031948 - 75034196	47,13	134 116	0.32/0.32/0.31	0.055/0.055/0.050	0.055/0.055/0.055
HMBS	2372	Chr. 11: 22521107 - 22523478	52,40	124 475	0.45/0.44/0.45	0.18/0.18/0.18	0.23/0.23/0.23
MYC	1794	Chr. 8: 42024572 - 42026365	49,39	3 191	5.5/5.3/5.3	6.6/6.4/6.2	3.5/3.2/3.0
SRC	1863	Chr. 20: 61680000 - 6169862	52,76	120 392	0.41/0.40/0.40	0.14/0.14/0.14	0.055/0.050/0.050
MxA	1882	Chr. 21: 28473387 - 28475268	49,15	4 631	3.4/3.2/2.5	1.1/1.1/0.8	2.6/2.4/1.9
PKR	1713	Chr. 2: 16170524 - 16172236	42,97	115 858	0.69/0.65/0.66	0.25/0.22/0.24	0.32/0.29/0.31
PML	2084	Chr. 15: 45116397 - 45118480	53,50	50 881	0.48/0.58/0.39	0.16/0.20/0.13	0.080/0.10/0.055
CXCL10	1577	Chr. 4: 1490756 - 1492332	37,67	103 739	0.44/0.44/0.45	0.16/0.16/0.17	0.055/0.055/0.050
CXCL13	1884	Chr. 4: 3074754 - 3076637	37,58	114 082	0.31/0.30/0.30	0.055/0.055/0.055	0.055/0.055/0.055
CCL17	2124	Chr. 16: 11062012 - 11064135	56,36	87 806	0.71/0.74/0.74	0.27/0.30/0.29	0.20/0.25/0.23
CCL20	1767	Chr. 2: 78889580 - 78891346	34,24	121 839	0.32/0.31/0.31	0.055/0.050/0.050	0.060/0.055/0.055
IL6	2070	Chr. 7: 22759130 - 22761199	45,17	118 519	0.41/0.45/0.41	0.13/0.13/0.12	0.11/0.15/0.11

D




Figure S8. (A) dHepaRG cells or PHH were infected with HBV, treated with IFN- α or LT β R-agonist BS1 for 10 days and 3D-PCR analysis was performed on a region of the *PrnP* gene. **(B)** 13 different amplicons from different genomic DNA regions from cells treated with IFN- α or BS1 were submitted for deep sequencing. The overall % of base exchanged for each base was analyzed. **(C)** The table gives the genes analyzed, length and genomic position of the amplicons, the coverage rate as well as mutations rates. **(D)** dHepaRG cells or primary human hepatocyte (PHH) were infected with HBV or mock-infected and treated with IFN- α or BS1 4 days later for 10 days. Total DNA was extracted and AP sites were quantified. Values given are the mean +/- standard deviation of biological triplicates from one experiment.

A

day 4

day 11




B**B****C****Nucleic acid binding ***

Figure S9. dHepaRG cells were infected with HBV and treated with LT β R-agonist BS1. **(A)** Affymetrix gene expression array analysis was confirmed by qRT-PCR analysis of the 50 most strongly up- and down-regulated genes. Results are shown as level of regulation according to the scale bars. **(B, C)** Regulated genes were classified according to their activity and binding properties. Stars indicate proteins for which nucleic acid binding properties have been described.




A**B****C**

Figure S10. (A) Indicated cells were infected with HBV and treated with LT β R-agonist BS1 (dHepaRG, dHepaRG-IRF3DN: 10 days; PHH: 7 days; PHH in uPA/SCID mice: 2 weeks). Expression of APOBEC3 family members were analyzed by qRT-PCR and values are shown as mean +/- standard deviation of independent replicates. (B) PHH were infected with HBV and treated with BS1 at 7 dpi for 10 days. Levels of cccDNA were compared to untreated PHH of the same donor (mock). (C) PCR analysis to detect the wild-type A3B allele or a common deletion of the A3B allele in genomic DNA extracted from PHH derived from different donors as well as from HepaRG cells.






Figure S11. dHepaRG cells were infected with HBV and treated with LT β R-agonist BS1. Expression levels of APOBEC3B and APOBEC3G after 10 days or at indicated time points were analyzed by qRT-PCR. Values show mean +/- standard deviation of independent replicates and data and were analyzed by t-test. * p<0.05, ** p<0.01 and *** p<0.001.

A



B



C

humans treated with PEG-IFN- α





Figure S12. (A) Indicated cells were infected with HBV and treated with IFN- α for 6 hours. Expression of APOBEC3 family members was analyzed by qRT-PCR and values are shown as mean \pm standard deviation of independent replicates. **(B)** Chimpanzees X0101 and X0233 were treated with 10 million IU of recombinant human IFN α -2a. Liver biopsies were obtained 8 hrs post treatment and expression of APOBEC3 genes were analyzed. **(C)** RNA isolated from livers of HCV-patients (n=3 to 6 at different time points) before and after treatment with 1.5 μ g/kg pegIFN α -2b were analyzed for the expression of APOBEC3 family members by gene expression array at indicated time points. Intensity values are given for the different APOBEC3 family members (upper graph) and fold changes were calculated when intensity values were above the threshold (lower graph).




Figure S13. (A) dHepaRG cells or (B) PHH were infected with HBV and treated with IFN- α . Expression of the indicated APOBEC3 mRNA after 6h or at the indicated time point were analyzed by qRT-PCR. Values show mean +/- standard deviation of independent replicates.





A**B****C****D**

Figure S14. dHepaRG cells were treated with IFN- α or mock-treated and fixed three days later. A3A, A3B, A3F or A3G localization were determined by **(A, C, D)** immunofluorescence staining or **(B)** Western blot analysis.






Figure S15. HBV-infected dHepaRG-tA-Vif cells were treated with **(A)** IFN- α or **(B, C)** LT β R-agonist BS1 for 10 days. 24h prior to HBV infection, HIV-Vif expression has been induced by doxycycline (dox). **(A, B)** Western blot analysis and **(C)** cccDNA qPCR were performed at the end of treatment. Values show the mean +/- standard deviation of independent replicates and data were analyzed by t-test. ** p<0.01. **(D)** A3A and **(E)** A3B were overexpressed in HepG2-H1.3 cells. 3D-PCR experiments were performed 5 days later and the resulting PCR products were cloned and sequenced. Content of each base is given.

A



B



C




Figure S16. dHepaRG-pEpi-H1.3 cells were treated with IFN- α (500 IU/ml) or BS1 (0.5 μ g/ml) for 11 days. (A) pEpi-H1.3 DNA, cccDNA and total intracellular HBV DNA were quantified by specific qPCR. (B) 3D-PCR specific for pEPIH.13 or cccDNA are shown. (C) HBV cccDNA 3D-PCR products obtained at indicated temperatures were cloned, sequenced and numbers of G/A mutations were evaluated.





Figure S17. HBV-infected dHepaRG and PHH were treated with IFN- α at 7 dpi for 3 days. A3A and HBC were analyzed by immunofluorescence staining and quantification (mean \pm SD) of A3A/HBC colocalisation in the nuclei from 5 different vision fields are given.

A



B





Figure S18. HepG2H1.3-A3A cells were analyzed by chromatin immunoprecipitation with anti-A3A, anti-HBc or using protein A/G beads only (negative control) and by qPCR using selective primers for (A) cccDNA (B) GAPDH, tumor suppressor p53, cellular/sarcoma tyrosine kinase (Src) and myelocytomatosis (c-myc).

A

A3A-transfected HepG2H1.3



B

A3A-Flag-transfected HuH7




Figure S19. (A) HepG2H1.3 cells were transfected with a plasmid expressing A3A. (B) Huh-7 cells were co-transfected with a plasmid expressing A3A-Flag and one containing a 1.1-fold HBV genome expressing all HBV genes. After 72h, cells were lysed and the indicated immunoprecipitations (IP) followed by Western blot for the A3A, Flag tag and HBV core (HBc) were performed.




Figure S20. Proximity Ligation Assay (PLA) staining in HepaRG cells with antibodies against HBc and A3A. Imaging was done by spinning disc microscopy and single cells were identified based on DAPI staining. PLA spots were automatically counted with a fixed threshold in all measurements.



A**B**

Figure S21. **(A)** Interaction of serial HBV Core (HBc)-YFP deletion mutants with A3G-CFP assessed by FACS-FRET in Huh7.5 hepatoma cells. Indicated are percentages of cells scoring FRET positive for each co-transfection. CFP-only and YFP-only as well as co-transfection of a highly membrane incorporated MEM fusion protein served as negative controls. A CFP-YFP fusion protein was used as positive control. Mean values and standard deviations were calculated from three to four independent experiments. **(B)** The potential interaction sites of modeled A3A (red) and HBV core protein (blue) are depicted: amino acids of HBV core locating within a distance of 3.1 Å from A3A, which are mainly located at residues 110 to 149, are highlighted in green indicating interaction. The red ball indicates the zinc atom embedded in the catalytic site of A3A. Our model indicates an interaction of HBV core α -helix 4 with A3A.

Table S1: Treatment of humanized uPA/SCID mice with LT β R agonists.

mouse*	treatment	total intracellular HBV DNA (copies/ cell)	cccDNA (copies/cell)
1	hu-IgG	570	1,79
2	hu-IgG	599	3,51
3	CBE11	84	0,84
4	CBE11	105	0,96

*Humanized uPA/SCID mice were infected with HBV and treated with chlodronate two times (day 0 and 5) to deplete murine liver macrophages and avoid Fc-receptor mediated killing of transplanted human hepatocytes. At day three, mice were treated either with LT β R agonistic antibody CBE11 or with control human IgG. Mice were sacrificed at day 10 and HBV rc and cccDNA in livers were analyzed.